arXiv:1302.1177v1 [gr-qc] 31 Jan 2013

Reheating temperature in non-minimal derivative coupling model

H. Mohseni Sadjadi *and Parviz Goodarzi
Department of Physics, University of Tehran,
P. O. B. 14395-547, Tehran 14399-55961, Iran

February 6, 2013

Abstract

We consider the inflaton as a scalar field described by a non-minimal derivative coupling model with a power law potential. We study the slow roll inflation, the rapid oscillation phase, the radiation dominated and the recombination eras respectively, and estimate e-folds numbers during these epochs. Using these results we determine the reheating temperature in terms of the spectral index and the amplitude of the power spectrum of scalar perturbations.

1 Introduction

To solve some dilemmas in the standard model of cosmology such as the flatness, the horizon, the monopoles problems and so on, inflation as an accelerated expansion era in the early universe was introduced by [1]. This scenario is now dubbed as old model of inflation, in which the universe underwent a de-Sitter expansion in a supercooled unstable false vacuum. Afterwards, by proposing a scalar field (inflaton) as the source of inflation, a new inflationary model was introduced in [2]. In this context, inflation was driven by the inflaton which slowly rolled down towards the minimum of its effective potential. To provide enough e-folds number, the potential must be nearly flat near its minimum.

The nature of this scalar field has not yet been identified, but a simple possible candidate might be the Higgs boson [3]. To adapt the inflaton to the Higgs boson, a non-minimally derivative coupling model in which the kinetic term of the inflaton is coupled to the Einstein tensor, was proposed in [4]. This model does not suffer from unitary violation problem and is safe from quantum corrections, besides slow roll inflation can be described by

*mohsenisad@ut.ac.ir

steep potentials in this framework[5, 6]. More general non-minimal derivative coupling model has also been considered in the literature to study the accelerated expansion of the universe in the early universe as well as the in the late time [7].

After the end of the inflation, the universe was cold and dominated by the inflaton scalar field energy. This energy had to be converted to relativistic particles to reheat the universe via a procedure called the reheating process [8]. A proposal for reheating, is the decay of the inflaton to ultra-relativistic particles (radiation) during a rapid coherent oscillation phase about the minimum of the potential. These particles interacted rapidly to become in thermal equilibrium characterized by the reheating temperature T_{reh}, and the universe entered on the radiation dominated era.

Although the exact value of T_{reh} has not yet been known, but some upper and lower bounds for this temperature have been obtained in the literature. By considering that the reheating process occurred before the big bang nucleosynthesis (BBN), and by combining constraints on light elements abundance and data obtained from large scale structure and cosmic microwave background (CMB), one can find a lower bound for T_{reh}, $4MeV \lesssim T_{reh}$ [9]. An upper bound may be taken as the energy scale at the end of inflation which is around the GUT scale $T_{reh} \lesssim 10^{16}GeV$. These assumptions give a wide range for the reheating temperature. In [10] by involving supersymmetry and considering the gravitino production, and on the base of cosmic microwave background (CMB) radiation data, this range was tightened to $6TeV \lesssim T_{reh} \lesssim 10^4 TeV$.

A more accurate method to determine T_{reh} in terms of CMB data was introduced in [11]. This method is based on determining the number of e-folds during the evolution of the universe from the inflation until the present time. Although in this context T_{reh} may be determined in terms of spectral index and amplitude of power spectrum of scalar perturbations, but due to uncertainties of theses quantities in WMAP7 data [12], a large relative uncertainty for T_{reh} is arisen: $\frac{\sigma(T_{reh})}{T_{reh}} \approx 53$, where $T_{reh} = 3.5 \times 10^6 GeV$.

In this paper we assume that the inflaton is a scalar field described by non-minimal derivative coupling model introduced in [4]. We study the inflationary era, the rapid oscillation phase of the inflaton, the radiation dominated epoch respectively and employ the method proposed in [11] to determine the reheating temperature in terms of the spectral index and power spectrum of scalar perturbations. Finally, the (range of the) value of T_{reh} is fixed via WMAP7 data.

We use units $\hbar = c = 1$ through the paper.

2 Evolution of the universe and the reheating temperature

We consider the spatially flat Friedmann-Lemaître- Robertson-Walker (FLRW) space-time

$$ds^2 = -dt^2 + a^2(t)(dx^2 + dy^2 + dz^2), \tag{1}$$

and choose an arbitrary length scale, λ_0, crossing the Hubble radius $R_H := \frac{1}{H} = \frac{a}{\dot{a}}$ at some time, denoted by t_*, during the inflation [13]. By using the red-shift relation

$$\frac{\lambda(t_*)}{\lambda_0} = \frac{a(t_*)}{a_0}, \tag{2}$$

where subscript "0" denotes the present time and by taking $a_0 = 1$, we obtain

$$\lambda_0 = \frac{1}{a(t_*)H(t_*)}. \tag{3}$$

This reference time will be used in division of the evolution of the universe into four parts as follows:

I- From t_* until the end of slow roll, denoted by t_e.

II- From t_e until the reheating or beginning of the radiation dominated epoch, denoted by t_{reh}.

III- From t_{reh} until recombination era, denoted by t_{rec}.

IV- From t_{rec} until the present time t_0.

The number of e-folds from t_* until t_0 is then given by

$$\begin{aligned} \mathcal{N} = \ln\left(\frac{a_0}{a_*}\right) &= \ln\left((\frac{a_0}{a_{rec}})(\frac{a_{rec}}{a_{reh}})(\frac{a_{reh}}{a_e})(\frac{a_e}{a_*})\right) \\ &= \mathcal{N}_{IV} + \mathcal{N}_{III} + \mathcal{N}_{II} + \mathcal{N}_I, \end{aligned} \tag{4}$$

where the subscripts denote the value of the parameter at their corresponding times.

In the following we will try to use eq.(4) to determine the reheating temperature in a non-minimal derivative coupling model in which the inflaton kinetic term is non-minimally coupled to Einstein tensor. This inflationary model is described by the action [4]

$$S_\phi = \int d^4x\sqrt{-g}\left[\frac{M_P^2}{2}R - \frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi + \frac{1}{2M^2}G^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi)\right], \tag{5}$$

where $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2}g^{\mu\nu}R$ is the Einstein tensor, R is the scalar curvature, M_P is the reduced planck mass given by $M_P = \sqrt{\frac{1}{8\pi G}} = 2.4 \times 10^{18} GeV$, and M is a scale with mass dimension.

2.1 Slow roll

In the era (I), the universe is dominated by the inflaton scalar field. The Friedmann equation is

$$H^2 = \frac{1}{3M_P^2}\rho_\phi, \tag{6}$$

where

$$\rho_\phi = \frac{1}{2}\left(1 + 9\frac{H^2}{M^2}\right)\dot{\phi}^2 + V(\phi), \tag{7}$$

is the energy and the upper dot is "$\dot{\cdot}$" $= \frac{d}{dt}$. The pressure is obtained as

$$P_\phi = \frac{1}{2}\left(1 - 3\frac{H^2}{M^2}\right)\dot{\phi}^2 - V(\phi) - \frac{1}{M^2}\frac{d(H^2\dot{\phi}^2)^2}{dt}. \tag{8}$$

With the help of continuity equation

$$\dot{\rho_\phi} + 3H(P_\phi + \rho_\phi) = 0, \tag{9}$$

one can derive the equation of motion of the inflaton as

$$\left(1 + \frac{3H^2}{M^2}\right)\ddot{\phi} + 3H\left(1 + \frac{3H^2}{M^2} + \frac{2\dot{H}}{M^2}\right)\dot{\phi} + V'(\phi) = 0. \tag{10}$$

In the continue, we restrict ourselves to high friction regime [4]

$$\frac{H^2}{M^2} \gg 1. \tag{11}$$

This choice, as we will see, by enhancing the slow roll procedure, enables us to consider more general steep potentials. During the slow roll, we have $\frac{H^2}{M^2}\dot{\phi}^2 \ll V(\phi)$, $\ddot{\phi} \ll H\dot{\phi}$, and therefore the Friedmann equation and inflaton equation of motion reduce to

$$\begin{aligned} H^2 &\approx \frac{1}{3M_P^2}V(\phi), \\ \dot{\phi} &\approx -\frac{M^2V'(\phi)}{9H^3} \end{aligned} \tag{12}$$

respectively. The slow roll parameters ϵ, δ satisfy

$$\epsilon := -\frac{\dot{H}}{H^2} \simeq \frac{M_P^2}{2}\frac{M^2}{3H^2}\frac{V'^2(\phi)}{V^2(\phi)} \ll 1,\ \delta := \frac{\ddot{\phi}}{H\dot{\phi}} \ll 1. \tag{13}$$

δ can be written as

$$\delta \simeq -\eta + 3\epsilon \tag{14}$$

where η is defined with

$$\eta := \frac{M_P^2}{3}\frac{M^2}{H^2}\frac{V''(\phi)}{V(\phi)}. \tag{15}$$

Eqs.(13-15), imply that by choosing an appropriate M in high friction regime (11), slow roll conditions do not oblige us to adopt approximately flat potentials.

Hereafter we will restrict ourselves to power law potential

$$V(\phi) = v\phi^n, \tag{16}$$

where v is a real number, and n is an even positive integer to guarantee that the potential has a minimum, about which the rapid oscillation of the inflaton occurs after slow roll.

The number of e-folds in the era (I) is

$$\begin{aligned}\mathcal{N}_I = \ln\left(\frac{a_e}{a_*}\right) = \int_{t_*}^{t_e} H dt &= \frac{1}{M_P^4 M^2}\int_{\phi_e}^{\phi_*}\frac{V^2(\phi)}{V'(\phi)}d\phi \\ &\simeq \frac{v}{n(n+2)M^2M_P^4}\phi_*^{n+2}.\end{aligned} \tag{17}$$

To obtain the above equation we have used (16), and $\phi_* \gg \phi_e$. To estimate ϕ_*, we consider the spectral index n_s [5],

$$n_s - 1 \simeq -2\epsilon - 2\delta \approx \frac{M_P^2M^2}{H_*^2}\left[-\frac{4}{3}\frac{V'^2(\phi_*)}{V^2(\phi_*)} + \frac{2}{3}\frac{V''(\phi_*)}{V(\phi_*)}\right]. \tag{18}$$

For the power law potential, this equation reduces to

$$1 - n_s = \frac{2M_P^4M^2n(n+1)}{v}\phi_*^{-(n+2)}. \tag{19}$$

The number od e-folds in the time interval I is then obtained by substituting ϕ_* from (19) into (17):

$$\mathcal{N}_I = 2\frac{(n+1)}{(n+2)(1-n_s)}. \tag{20}$$

2.2 Reheating era

At the end of slow roll we have $\epsilon(\phi_e) \simeq 1$, which yields

$$\phi_e^{n+2} = \frac{M_P^4M^2n^2}{2v}. \tag{21}$$

At this time the energy density is approximately given by

$$\rho_e \simeq V(\phi_e) = v\left(\frac{M^2M_P^4n^2}{2v}\right)^{\frac{n}{n+2}}, \tag{22}$$



Figure 1: $\varphi := \frac{\phi}{M_P}$ in terms of dimensionless time $\tau = mt$, for $\frac{m^2}{M^2} = 10^8$ with initial conditions $\{\varphi(1) = 0.056,\ \dot{\varphi}(1) = 0\}$, for the quadratic potential.

and the scalar field commences a rapid coherent oscillation around the bottom of the potential (see fig.(1), depicted for the quadratic potential $V(\phi) = \frac{1}{2}m^2\phi^2$) via numerical methods. In this high frequency regime, where the behavior of the scalar field is opposite to the slow roll, the energy density and the Hubble parameter decrease very slowly (insignificantly) in one period of oscillation [14].

Now let us consider $\gamma = 1 + w_\phi$, where w_ϕ is the effective equation of state parameter of the the scalar field: $w_\phi = \frac{\langle P_\phi \rangle}{\langle \rho_\phi \rangle}$. $< ... >= \frac{\int_t^{t+T} ...dt'}{T}$ is the time average over an oscillation whose period is T. With the help of (7) and (8), γ reads

$$\begin{aligned} \gamma &= 1 + \frac{< P_\phi >}{< \rho_\phi >} \\ &= \frac{\left\langle \frac{3H^2}{M^2}\dot{\varphi}^2 - \frac{1}{M^2}\frac{d\left(H\dot{\varphi}^2\right)}{dt} \right\rangle}{\langle \rho_\phi \rangle}. \end{aligned} \tag{23}$$

Using $dt = \frac{1}{\dot{\phi}}d\phi$, (7), (16), and by noting that the Hubble parameter does not change considerably in one period, we obtain

$$\begin{aligned} \gamma &= \frac{2}{3}\frac{\langle \rho_\phi - V(\phi) \rangle}{\langle \rho_\phi \rangle} \\ &= \frac{2}{3\Phi^n}\frac{\int_{-\Phi}^{\Phi}\sqrt{\Phi^n - \phi^n}d\phi}{\int_{-\Phi}^{\Phi}\frac{d\phi}{\sqrt{\Phi^n - \phi^n}}} \\ &= \frac{2}{3}\frac{\int_0^1\sqrt{1-x^n}dx}{\int_0^1\frac{1}{\sqrt{1-x^n}}dx} = \frac{2n}{3n+6}. \end{aligned} \tag{24}$$

Φ is the amplitude of $\phi(t)$, i.e. $V(\Phi) \simeq \rho_\phi$. If the non-minimal derivative coupling was absent we would have $\gamma = \frac{2n}{n+1}$ [14].

We assume that in this epoch the inflaton decays to ultra-relativistic particles (whose the energy density is denoted by ρ_r), to reheat the universe. From the beginning of rapid oscillation, i.e. $\rho_r = 0$, until $\rho_r = \rho_{reh} \simeq \rho_\phi$, which is the beginning of radiation dominated era, the universe is approximately dominated by the rapidly oscillating scalar field. Therefore, in this era the Hubble parameter can be approximated by [15]

$$H \simeq \frac{2}{3\gamma t}, \tag{25}$$

where γ is given by (24). The scale factor during this era scales as

$$\frac{a_{reh}}{a_e} \simeq \left(\frac{\rho_e}{\rho_{reh}}\right)^{\frac{n+2}{2n}}. \tag{26}$$

At t_{reh} we can estimate the energy density as

$$\rho_{reh} \simeq \frac{g_{reh}}{30}\pi^2 T_{reh}^4, \tag{27}$$

where T_{reh} is the temperature of ultra relativistic particles at t_{reh}, and g_{reh} is the number of (massless) degrees of freedom corresponding to the ultra-relativistic particles present in the model at t_{reh} [16]. Collecting all together, we can estimate the number of e-folds during rapid oscillation

$$\begin{aligned} \mathcal{N}_{II} &= \ln\left(\frac{a_{reh}}{a_e}\right) \\ &= \ln\left(\sqrt{\frac{M_P^4 M^2 n^2}{2}}\left(\frac{30}{g_{reh}\pi^2 T_{reh}^4}\right)^{\frac{n+2}{2n}} v^{\frac{1}{n}}\right). \end{aligned} \tag{28}$$

To be more specific we must determine v. To do so, consider the power spectrum of the scalar perturbation

$$\mathcal{P} = \frac{H^2}{8\pi^2 M_P^2 \epsilon}, \tag{29}$$

which is computed at the horizon crossing $k = k_0 = a_* H_*$ (see (2) and (3)), where $k_0 = \frac{1}{\lambda_0}$ is a pivot scale. Thus

$$H_* = 2\pi M_P \sqrt{2\epsilon \mathcal{P}(k_0)}. \tag{30}$$

Using this equation together with (19) and $H_*^2 \simeq \frac{1}{3M_P^2} V(\phi_*)$, and after some computations we find that

$$v = \left(\frac{1-n_s}{1+n}\right)^{1+n} \left(6\pi^2 \mathcal{P}(k_0)\right)^{\frac{n+2}{2}} \left(2M^2\right)^{-\frac{n}{2}} M_P^4 n. \tag{31}$$

Substituting (31) into (28) yields

$$\mathcal{N}_{II} = \ln\left[\frac{1}{2}\left(\frac{n(1-n_s)}{1+n}\right)^{\frac{1+n}{n}}\left(\frac{180M_P^4\mathcal{P}(k_0)}{g_{reh}T_{reh}^4}\right)^{\frac{n+2}{2n}}\right]. \tag{32}$$

2.3 Radiation dominated and recombination eras

In the radiation dominated era the universe is constituted of ultra-relativistic particles which are in thermal equilibrium, and undergoes an adiabatic expansion during which the entropy per comoving volume is conserved: $dS = 0$ [16]. In this era the entropy density, defined by $s = Sa^{-3}$, is obtained as [16]

$$s = \frac{2\pi^2}{45}g_{reh}T^3. \tag{33}$$

So we have

$$\frac{a_{rec}}{a_{reh}} = \frac{T_{reh}}{T_{rec}}\left(\frac{g_{reh}}{g_{rec}}\right)^{\frac{1}{3}}. \tag{34}$$

In the recombination era, g_{rec} corresponds to photons degrees of freedom and consequently $g_{rec} = 2$. Hence

$$\mathcal{N}_{III} = \ln\left(\frac{T_{reh}}{T_{rec}}\left(\frac{g_{reh}}{2}\right)^{\frac{1}{3}}\right). \tag{35}$$

The temperature decreases by the expansion of the universe via $T(z) = T(z=0)(1+z)$, where z is the redshift parameter. Therefore we can express T_{rec} in terms of T_{CMB} as

$$T_{rec} = (1 + z_{rec})T_{CMB}. \tag{36}$$

We have also

$$\frac{a_0}{a_{rec}} = (1 + z_{rec}) \tag{37}$$

Thus

$$\mathcal{N}_{III} + \mathcal{N}_{IV} = \ln\left(\frac{T_{reh}}{T_{CMB}}\left(\frac{g_{reh}}{2}\right)^{\frac{1}{3}}\right). \tag{38}$$

2.4 Reheating temperature

So far, we have determined the number of e-folds in the the right hand side of (4). To obtain the reheating temperature we need also to determine $\mathcal{N}$ in (4). Taking $a_0 = 1$, the number of e-folds from the horizon crossing until the present time is derived as $\mathcal{N} = \exp(\Delta)$, where

$$\Delta = \frac{1}{a_*} = \frac{H_*}{k_0} = \frac{2\pi M_P}{k_0}\sqrt{\frac{2n(1-n_s)}{(1+n)}\mathcal{P}_s(k_0)}. \tag{39}$$

To obtain the above relation we have made use of (30) and

$$\epsilon \simeq \frac{n(1-n_s)}{4(n+1)}, \tag{40}$$

which is derived from (13-15) and (19).

Using (4), (20), (32), (38), and (39) we finally arrive at

$$T_{reh} = \alpha T_{CMB}^{-\frac{n}{n+4}} \tag{41}$$

where α is defined through

$$\begin{aligned} \alpha &= g_{reh}^{\frac{n+6}{6n+24}} M_P \left(\frac{k_0}{2^{\frac{11}{6}}\pi}\right)^{\frac{n}{n+4}} \left(\frac{180n(1-n_s)}{n+1}\right)^{\frac{n+2}{2n+8}} \times \\ &\exp\left(\frac{2n(n+1)}{(n+2)(n+4)(1-n_s)}\right) \mathcal{P}_s^{\frac{1}{n+4}}(k_0) \end{aligned} \tag{42}$$

α, up to this order of approximation, is independent of v and M. Adopting the pivot scale as $k_0 = 0.002 Mpc^{-1}$, $\mathcal{P}_s$ and n_s are measured as [12]

$$\begin{aligned} \mathcal{P}_s(k_0) &= 2.441^{+0.088}_{-0.092} \times 10^{-9} \\ n_s &= 0.963 \pm 0.012. \end{aligned} \tag{43}$$

For the quartic potential, $n = 4$, (41) reduces to

$$T_{reh} = 1.927 g_{reh}^{-\frac{5}{24}} M_P (1-n_s)^{\frac{3}{8}} \mathcal{P}_s^{\frac{1}{8}}(k_0) \exp\left(\frac{5}{6(1-n_s)}\right) \left(\frac{k_0}{T_{CMB}}\right)^{\frac{1}{2}}. \tag{44}$$

By taking $g_{reh} = 106.75$, which is the ultrarelativistic degrees of freedom at the electroweak energy scale, we find $T_{reh}\left(\mathcal{P}_s(k_0) = 2.441 \times 10^{-9}, n_s = 0.963\right) = 1.93 \times 10^{12} GeV$. Considering the uncertainties, one can find that the reheating temperature lies in the range

$$0.86 \times 10^{10} GeV \leq T_{reh} \leq 0.83 \times 10^{17} GeV. \tag{45}$$

For quadratic potential, $n = 2$, (41) reduces to

$$T_{reh} = 2.205 g_{reh}^{-\frac{2}{9}} M_P (1-n_s)^{\frac{1}{3}} \mathcal{P}_s^{\frac{1}{6}}(k_0) \exp\left(\frac{1}{2(1-n_s)}\right) \left(\frac{k_0}{T_{CMB}}\right)^{\frac{1}{3}}, \tag{46}$$

giving $T_{reh}\left(\mathcal{P}_s(k_0) = 2.441 \times 10^{-9}, n_s = 0.963, g_{reh} = 106.75\right) = 6.53 \times 10^{12} GeV$. By taking into accounts the uncertainties, we obtain

$$0.26 \times 10^{12} GeV \leq T_{reh} \leq 0.38 \times 10^{16} GeV. \tag{47}$$

In the minimal model, for the quadratic potential, the reheating temperature was obtained as [11]

$$T_{reh} = 0.017 M_P (1-n_s)^{\frac{1}{2}} \mathcal{P}_s^{-\frac{1}{2}}(k_0) \exp\left(\frac{6}{(1-n_s)}\right) \left(\frac{k_0}{T_{CMB}}\right)^{3}. \tag{48}$$

Note that in contrast to the non-minimal derivative coupling model, T_{reh} in (48) does not depend on relativistic degrees of freedom g_{reh}. (48) yields $T_{reh}\left(\mathcal{P}_s(k_0) = 2.441 \times 10^{-9}, n_s = 0.963\right) = 3.5 \times 10^6 GeV$. This temperature is very less than that we have obtained in our model at these values of the amplitude and the spectral index. But we must note that the uncertainty in the minimal model with quadratic potential which is given by [11]

$$\frac{\sigma(T_{reh})}{T_{reh}} \approx 53, \tag{49}$$

is very large with respect to the uncertainty in the non-minimal model which is obtained as

$$\frac{\sigma(T_{reh})}{T_{reh}} = 4.275. \tag{50}$$

3 Conclusion

Non-minimal derivative coupling model with a power law potential was employed to describe the inflation (see (5)). In this context, the slow roll inflationary phase of the inflaton was discussed in high friction regime (see (11)). The rapid oscillation phase after the slow roll, during which the inflaton decays to ultra-relativistic particles, was studied. From the beginning of this rapid oscillation until the radiation dominated epoch, the equation of state parameter of the universe can be approximated by a constant (see (24)), which enabled us to compute the number of e-folds in this era (see (28)). We also estimated the number of e-folds number in radiation dominated, and recombination era, in the same way as the minimal model (see (38)). By gathering all these results together, we obtained the reheating temperature in terms of T_{CMB}, spectral index and the amplitude of the power spectrum of scalar perturbations (see (41)). Finally according to WMAP 7 data, we determined the value of T_{reh} which is much bigger than the reheating temperature obtained in minimal model. This is due to high friction regime adopted in this paper, which enhances the decay of the inflaton. We also showed that the uncertainty in our result is very smaller with respect to the minimal model.

References

[1] A. H. Guth, Phys. Rev. D 23, 347(1981).

[2] A. Linde, Phys. Lett. B 129, 177 (1983); A.Linde, Particle Physics and Inflationary Cosmology (Harwood, Chur, Switzerland, 1990).

[3] R. N. Lerner and J. McDonald, Phys. Rev. D 82, 103525 (2010); R. N. Lerner and J. McDonald, Phys. Rev. D 83, 123522 (2011); K. Nakayama and F. Takahashi, arXiv:1108.3762 [hep-ph].

[4] C. Germani and A. Kehagias, Phys. Rev. Lett. 105, 011302 (2010).

[5] C. Germani and Y. Watanabe, JCAP 07(2011)031; C. Germani and A. Kehagias, arXiv:1003.4285 [astro-ph.CO].

[6] C. Germani and A. Kehagias, Phys. Rev. Lett. 106, 161302 (2011); C. Germani, arXiv:1112.1083v1 [astro-ph.CO].

[7] S. V. Sushkov, Phys. Rev. D 80, 103505 (2009); E. N. Saridakis and S. V. Sushkov, Phys. Rev. D 81, 083510 (2010); L. N. Granda, arXiv:1104.2253 [hep-th]; S. Tsujikawa, arXiv:1201.5926v1 [astro-ph.CO]; H. M. Sadjadi, Phys. Rev. D 83,107301 (2011); G. Gubitosi and E. V. Linder, Phys. Lett. B 703, 113 (2011); A. Banijamali and B. Fazlpour, Phys. Lett. B 703, 366 (2011); S. A. Appleby, A. De Felice, and E. V. Linder, JCAP 1210 (2012) 060, arXiv:1208.4163 [astro-ph.CO]; S. V. Sushkov, Phys. Rev. D85, 123520 (2012).

[8] A. Albresht, P. J. Steinhardt, M. S. Turner, and F. Wilczek, Phys. Rev. Lett. 48, 1437 (1982); L. Abbott, E. Farhi, and M. Wise, Phys. Lett. B 117, 29 (1982); A. Doglov and A.Linde, Phys. Lett. B 116, 329 (1982); L. Kofman, A. D. Linde and A.A. Starobinsky, Phys. Rev. Lett 73, 3195 (1994); L. Kofman, A. D. Linde and A.A. Starobinsky, Phys. Rev. D 56, 3258 (1997).

[9] S. Hannestad, Phys. Rev. D 70, 043506 (2004).

[10] J. Martin and C. Ringeval, Phys. Rev. D 82, 023511 (2010).

[11] J. Mielczarek, Phys. Rev. D 83, 023502 (2011).

[12] E. Komatsu et al., Astrophys. J. Suppl. 192, 18 (2011), arXiv:1001.4538 [astro-ph.CO].

[13] A. R. Liddle and D. H. Lyth, Cosmological Inflation and Large-Scale Structure (Cambridge University Press 2000); A. R. Liddle and D.H. Lyth, Phys. Rept. 231, 1 (1993).

[14] Y. Shtanov, J. Traschen, and R. Brandenberger, Phys. Rev. D 51, 5438 (1995).

[15] H. Mohseni Sadjadi and P. Goodarzi, arXiv:1203.1580v2 [gr-qc].

[16] E. Kolb and M. Turner, The Early Universe (Addison-Wesley Publishing Company, Redwood City, California, 1990).